



<input type="checkbox"/>	<b>Drug-induced thrombocytopenia</b> (15 ml EDTA-Blood und 5 ml Native blood)	
Please add 2 – 3 tablets (minimum 30mg) or 1 ampule of the suspected drug  Increase of platelet count after discontinuation of drug  yes <input type="checkbox"/> no <input type="checkbox"/>		<b>PLT:</b> <b>GPT/L</b> Drugs (started 14 days before occurrence of thrombocytopenia):
<input type="checkbox"/>	<b>Platelet – specific antibodies</b>	
<b>PLT:</b> <b>GPT/L</b>  Immune thrombocytopenia (ITP) <input type="checkbox"/> (15 ml EDTA-Blood and 5 ml Native blood)  Neonatal alloimmune thrombocytopenia (NAIT) <input type="checkbox"/> <b>father:</b> 20 ml EDTA - Blood (required) <b>mother:</b> 10 ml Native blood + 10 ml EDTA - Blood (required) <b>child:</b> 1 ml EDTA – Blood (optional) (cord blood possible)		Refractory platelet count after platelet transfusion <input type="checkbox"/> (5 ml EDTA-Blood and 10 ml Native blood)  Posttransfusion purpura (PTP) <input type="checkbox"/> (5 ml EDTA-Blood and 10 ml Native blood)
Does the patient receive blood transfusions (during the last weeks)? <b>yes</b> <input type="checkbox"/> (Date of the last transfusion:                      ) <b>no</b> <input type="checkbox"/>		
Pregnancy or miscarriage? <b>yes</b> <input type="checkbox"/> <b>no</b> <input type="checkbox"/>		
<input type="checkbox"/>	<b>Inherited platelet disorders</b> 5-10 ml EDTA-blood; (3-5 ml EDTA-blood, children), 10 air-dried unstained blood smears, 2 stained blood smears (See also request form: <a href="http://www.medizin.uni-greifswald.de/transfus/index.php?id=389">http://www.medizin.uni-greifswald.de/transfus/index.php?id=389</a> )	
<b>Screening</b>	<input type="checkbox"/> thrombocytopenia <input type="checkbox"/> platelet function disorder	
<b>Bleeding tendency</b> <input type="checkbox"/> mild <input type="checkbox"/> intermediate <input type="checkbox"/> severe	please describe (e.g. easy bruising, menorrhagia, joint bleeding, etc.)	
<input type="checkbox"/>	<b>TRALI, neutrophil antibodies</b>	
<b>Blood donor:</b> 5 ml EDTA-Blood und 10 ml Native blood + <b>Patient:</b> 10 ml EDTA-Blood and 10 ml Native blood		
Transfused blood product	Red blood cell-concentrate <input type="checkbox"/> platelet concentrate <input type="checkbox"/> plasma <input type="checkbox"/> others: _____	

**Children:** Please contact our staff for reduced blood volumes: [thrombolab@uni-greifswald.de](mailto:thrombolab@uni-greifswald.de)

**Contact person:**  
 Thrombozytenlabor: Dr. K. Althaus/Dr. G. Hron (Thrombozytenlabor: 0049-3834 – 86-5475)

Place/Date/Seal:

Signature: